



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Entire genomic sequence of novel canine papillomavirus type 13**

Lange, C E ; Ackermann, M ; Favrot, C ; Tobler, K

**Abstract:** Papillomaviruses are associated with benign and malignant neoplasias of the skin and mucous membranes. The sequence of a novel canine papillomavirus was determined from DNA detected in the oral cavity of a dog. The sequence of the novel virus canine papillomavirus type 13 (CPV13) shares the highest levels of similarity with the Tau papillomaviruses CPV2 and CPV7.

DOI: <https://doi.org/10.1128/JVI.01553-12>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-65931>

Journal Article

Accepted Version

Originally published at:

Lange, C E; Ackermann, M; Favrot, C; Tobler, K (2012). Entire genomic sequence of novel canine papillomavirus type 13. *Journal of Virology*, 86(18):10226-10227.

DOI: <https://doi.org/10.1128/JVI.01553-12>

# **Entire genomic sequence of novel canine papillomavirus type 13**

Christian E. Lange,<sup>a,b</sup> Mathias Ackermann,<sup>b</sup> Claude Favrot<sup>a</sup> and Kurt Tobler<sup>b</sup>

Dermatology Department, Clinic for Small Animal internal Medicine, Vetsuisse Faculty,  
Zurich, Switzerland,<sup>a</sup> and Institute of Virology, Vetsuisse Faculty, Zurich, Switzerland<sup>b</sup>

Papillomaviruses are associated with benign and malignant neoplasias of the skin and mucous membranes. The sequence of a novel canine papillomavirus was determined from DNA detected in the oral cavity of a dog. The sequence of the novel virus Canine Papillomavirus 13 (CPV13) shares the highest level of similarity with the Tau papillomaviruses CPV2 and CPV7.

Papillomaviruses are non-enveloped, icosahedral particles, approximately 50 nm in diameter, with a circular, double stranded DNA genome of about 8.000 base pairs (bp). Typically, they are host species-specific and tissue-restricted putative pathogens. Many of the known papillomaviruses are associated with benign and malignant neoplasias of the keratinizing and non-keratinizing skin in humans and animals (3). More than 150 human and also many animal papillomaviruses types have been discovered, illustrating a broad genetic diversity (1, 3, 4). A dozen complete and a handful of partial canine papillomavirus (CPV) sequences have been published and linked to various neoplasias. Thus far, all CPVs were allocated to three different papillomavirus genera, i.e. Lambda, Tau or Chi (1, 2, 5-14).

A cytobrush sample of a mixed breed dog showing symptoms of oral papillomatosis was taken for diagnostic purposes. Thereof, total DNA was isolated, a circular DNA amplified by rolling circle amplification (RCA), and partially as well as entirely cloned into the BamHI,

24 ClaI or EagI site of a pBluescript II KS+ vector (Stratagene). The sequences of the RCA  
25 product and the genomic clones were determined independently using an ABI 377 (Applied  
26 Biosystems) sequencer and primary sequences were assembled using Contigexpress software  
27 (Vector NTI Informax, Invitrogen), revealing a papillomavirus genome of 8228 bp with a GC  
28 content of 47%. Pairwise sequence alignments were performed with the obtained sequence  
29 using the Needleman-Wunsch algorithm, and a phylogenetic tree was calculated based on the  
30 coding sequences for the E6, E7, E1, E2, L2 and L1 proteins (not shown) (7).

31 The characteristic open reading frames E1, E2, E4, E6, E7, L1 and L2 as well as two non-  
32 coding regions between L1 and E6 (478 bp) and between E2 and L2 (914 bp) were identified  
33 on the nucleotide sequence of the novel isolate. Dyad symmetry repeats  
34 (TTGTTGTAAACAACAA) in a modified form flanked by E2 binding sites (ACC-N<sub>6</sub>-GGT)  
35 were located around hundred nucleotides upstream of E6 putatively marking the origin of  
36 replication. Polyadenylation signals (AATAAA) were found near the 5' end of the L2 as well  
37 as in the terminal third and at the 3' end of the L1 open reading frame. A typical pRB-binding  
38 motif (LXCXE) in the E7 amino acid sequence of the novel isolate was not detected, an  
39 absence shared with the Tau papillomaviruses CPV2 and CPV7.

40 The L1 nucleotide sequence shared 63% identity with CPV7 and 62% with CPV2, while it  
41 shared less than 60% with all other published papillomavirus L1 sequences. The highest  
42 degree of similarity in the E1 amino acid sequence was found with CPV7 (68.8%) and CPV2  
43 (68.4). On the level of E6 and E7 the highest similarity was observed with CPV2 (55.3% and  
44 68.4%) and CPV7 (50.4% and 64.3). Upon the phylogenetic analysis the novel papillomavirus  
45 sequence clustered with the two previously described Tau CPVs. Based on the overall  
46 analyses the new isolate was designated as CPV13.

47 Taken together the findings suggest that CPV13 might be the third member of the  
48 papillomavirus genus Tau putatively establishing a new species within it.

49 **Nucleotide sequence accession number.** The nucleotide sequence data of CPV13 were  
50 deposited in GenBank under accession no. JX141478.

51

## 52 **Acknowledgements**

53 The authors thank Katja B. Egli, Anita Meier, Romy Schleiss and Oliver R. Wittwer all University of  
54 Zurich, as well as Elisabeth Vetsch and Mario Wickert of the Institute of Virology, Vetsuisse Faculty  
55 Zurich for support with the clinical or labwork. This work was self financed.

## 56 **References**

- 57 1. Bernard, H., R. Burk, Z. Chen, K. van Doorslaer, H. Hausen & E. de Villiers. 2010.  
58 Classification of papillomaviruses (PVs) based on 189 PV types and proposal of  
59 taxonomic amendments. *Virology*, 401, 70-9.
- 60 2. Delius H, Van Ranst M, Jenson A, zur Hausen H, Sundberg J. 1994. Canine oral  
61 papillomavirus genomic sequence: a unique 1.5-kb intervening sequence between the E2  
62 and L2 open reading frames. *Virology*, 204, 447-52.
- 63 3. Howley PM & Lowy DR. 2007. Papillomaviruses. In: D. M. Knipe & P. M. Howley  
64 (Eds.), *Fields Virology*, Lippincott Williams & Wilkins, Philadelphia, pp. 2299-2354.
- 65 4. Kovanda A, Kocjan BJ, Luzar B, Bravo IG, Poljak M. 2011. Characterisation of novel  
66 cutaneous human papillomavirus genotypes HPV-150 and HPV-151. *PLoS One* 6,  
67 e22529.
- 68 5. Lange CE, Tobler K, Ackermann M., Panakova L, Thoday KL, Favrot C. 2009. Three  
69 novel canine papillomaviruses support taxonomic clade formation. *J Gen Virology* 90,  
70 2615-2621.

- 71 6. Lange CE, Tobler K, Brandes K, Breithardt K, Ordeix L, Von Bomhard W, Favrot C.  
72 2010. Canine inverted papillomas associated with DNA of four different  
73 papillomaviruses. *Vet Dermatol* 21, 287-91.
- 74 7. Lange CE, Tobler K, Lehner A, Vetsch E, Favrot C. 2012. A case of a canine pigmented  
75 plaque associated with the presence of a Chi-papillomavirus. *Veterinary Dermatology* 23,  
76 76-e19.
- 77 8. Luff JA, Affolter VK, Yeargan B, Moore, PF. 2012. Detection of six novel  
78 papillomavirus sequences within canine pigmented plaques. *J Vet Diagn Invest*
- 79 9. Munday JS, O'Connor KI, 2011. Smits B. Development of multiple pigmented viral  
80 plaques and squamous cell carcinoma in a dog infected by a novel papillomavirus. *Vet*  
81 *Dermatol* 22, 104-10.
- 82 10. Tobler K, Favrot C, Nespeca G, Ackermann M. 2006. Detection of the prototype of a  
83 potential novel genus in the family Papillomaviridae in association with canine  
84 epidermodysplasia verruciformis. *J Gen Virol*, 87, 3551-7.
- 85 11. Tobler K, Lange C, Carlotti DN, Ackermann M, Favrot C. 2008. Detection of a novel  
86 papillomavirus in pigmented plaques of four pugs. *Veterinary Dermatology* 19, 21-25.
- 87 12. Yuan H, Ghim S, Newsome J, Apolinario T, Olcese V, Martin M, Delius H, Felsburg P,  
88 Jenson B, Schlegel R. 2007. An epidermotropic canine papillomavirus with malignant  
89 potential contains an E5 gene and establishes a unique genus. *Virology*, 359, 28-36.
- 90 13. Yuan H, Luff J, Zhou D, Wang J, Affolter V, Moore P, Schlegel R. 2012. Complete  
91 Genome Sequence of Canine Papillomavirus Type 9. *J. Virol.* 86, 5966.
- 92 14. Zaugg N, Nespeca G, Hauser B, Ackermann M, Favrot C. 2005. Detection of novel  
93 papillomaviruses in canine mucosal, cutaneous and in situ squamous cell carcinomas. *Vet*  
94 *Dermatol*, 16, 290-8.